

## PART II: HIV-1 CTL EPITOPES

### SUMMARY

Part II includes tables, maps, and associated references of HIV-specific CTL epitopes arranged sequentially according to the location of the proteins in the HIV-1 genome. We attempted to make this section as comprehensive as possible, requiring that the epitope be contained within a defined region of a maximum of 30 amino acids, but not that the precise boundaries be defined. Studies that were based on the analysis of whole proteins are described at the end of each protein section. The same epitope can have multiple entries, and each entry represents a single publication in this section of the database. For more recent updates and useful searching capabilities, please see our web site: <http://hiv-web.lanl.gov/immunology>. For a concise listing of the best defined CTL epitopes, see the summary by Christian Brander and Philip Goulder in Part I of this compendium. CTL protein reactions with no well-defined epitopes are listed at the end of each protein section.

### A. CTL EPITOPE TABLES

Each CTL reference has a six part basic entry:

- **HXB2 Location:** The viral strain HXB2 (GenBank Accession Number K03455) is used as a reference strain throughout this publication. The position of the defined epitope location on the sequence of the HXB2 protein is indicated. Obviously HXB2 may not be identical to a given defined reactive sequence, so we are simply indicating the location of the aligned positions. The HXB2 numbering is used in the protein maps in this database and is the reference strain in the HIV Sequence Compendium. HXB2 was chosen as the reference clone because it is the most intensively studied strain in terms of immunogenicity, structure, and function.
- **Author Location:** The amino acid positions of the epitope boundaries and the reference sequence are listed as given in the primary publication. Frequently, these positions as published are imprecise, and do not truly correspond to the numbering of the sequence, but they provide a reasonable guide to the peptide's approximate location in the protein. Also, in many cases the reference sequence identification was not provided, and in such cases it is not possible to use these numbers to specify precise

locations. If you are interested in finding the precise positions of epitopes you are studying relative to the HXB2 strain, please try using the interactive position locator at our web site: <http://hiv-web.lanl.gov/NUM-HXB2/HXB2.MAIN.html>.

- **Epitope Sequence:** The amino acid sequence of the epitope of interest as defined in the reference, based on the reference strain used in the study defining the epitope. On rare occasions, when only the epitope location and not the actual epitope sequence was specified in the original publication, and the sequences were numbered inaccurately by the primary authors, we may have misrepresented the epitope's amino acid sequence. Therefore epitopes that were not explicitly written out in the text in the primary publication, those that we determined by looking up the reference strain and the numbered location, are followed by a question mark in the table.
- **Immunogen:** The antigenic stimulus of the CTL response.
- **Species(HLA):** The species responding and HLA of MHC specificity of the epitope.
- **Reference:** The primary reference (sometimes two or more directly related studies are included).

Following the entry for a given CTL epitope are brief comments explaining the context in which the epitope was studied. If the same epitope was studied in several labs, each study is cited in its own bulleted entry.

### B. HIV PROTEIN EPITOPE MAPS

Because of the increasing number of defined epitopes, only HIV CTL epitopes and mapped to within a region of 21 amino acids or less, with a known HLA specificity, are indicated on the HIV protein epitope maps.

The location and HLA restriction elements of CTL epitopes are indicated on protein sequences of HXB2. These maps are meant to provide the relative location of epitopes on a given protein, but the HXB2 sequence may not actually carry the epitope of interest, as it may vary relative to the sequence for which the epitope was defined. Epitopes are numbered in bold on the maps; the map numbering corresponding to the numbering of the epitope sequence alignments.

## **HIV CTL Epitopes**

### **C. REFERENCES AND NOTES**

#### **ALIGNMENTS**

To conserve space, no epitope alignments are provided in this book, but they can be generated using the CTL epitope search tool at

<http://hiv-web.lanl.gov/immunology>